## **Food Allergy Action Plan**

Student's Name:D.O.B:				Place Child's	
ALLERGY TO:			Picture Here		
Asthmatic Yes* No *Higher risk fo	or severe reaction			Here	
♦ <u>STEP 1: TREATMENT</u> ♦					
Symptoms:		Give Checked Medication**:  To be determined		determined	
If a food allergen has been ingested, but <i>no symptoms</i> :		□ EpiPen	☐ Antihistam	autnorizing	
<ul> <li>Mouth Itching, tingling, or swelling of lips, tongue, mouth</li> </ul>		□ EpiPen	☐ Antihistamine ☐		
• Skin Hives, itchy rash, swelling of the face or extremities		☐ EpiPen	☐ Antihistamine		
<ul> <li>Gut Nausea, abdominal cramps, vomiting, diarrhea</li> </ul>		☐ EpiPen	☐ Antihistamine		
■ Throat + Tightening of throat, hoarseness, hacking cough		☐ EpiPen	☐ Antihistamine		
■ Lung + Shortness of breath, repetitive coughing, wheezing		☐ EpiPen	☐ Antihistamine		
■ Heart + Thready pulse, low blood pressure, fainting, pale, blueness		☐ EpiPen	☐ Antihistamine		
• Other +		☐ EpiPen	☐ Antihistamine		
■ If reaction is progressing (several of the above areas	☐ EpiPen	☐ Antihistamine			
The severity of symptoms can quickly change. † Potentially life-threatening.					
<b>DOSAGE Epinephrine:</b> inject intramuscularly (circle one) EpiPen EpiPen Jr. (see reverse side for instructions)					
Antihistamine: give					
Other: give					
medication/dose/route					
◆ <u>STEP 2: EMERGENCY CALLS</u> ◆					
1. Call 911 (or Rescue Squad:). State that an allergic reaction has been treated, and additional epinephrine may be needed)					
2. Dr at					
3. Emergency contacts: Name/Relationship Phone Number(s)					
a 1.)		2.)			
b 1.)		2.)			
c 1.)		2.)			
EVEN IF PARENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICATE OR TAKE CHILD TO MEDICAL FACILITY!					
Parent/Guardian Signature			Date		

Date\_

Doctor's Signature\_

(Required)